

Lignans and chronic diseases: results from the PREDIMED trial

Tresserra-Rimbau A^{1,2*}, Estruch R^{2,3}, Martínez-González MA^{2,4,5}, Fitó M^{2,6}, Corella D^{2,7}, Salas-Salvadó J^{2,8}, Guasch-Ferré M¹⁰, Rimm EB^{9,10}, Lamuela-Raventós RM^{1,2}, on behalf of the PREDIMED Study Investigators

¹Department of Nutrition, Food Science and Gastronomy, XaRTA, INSA-UB, School of Pharmacy and Food Science, University of Barcelona, Barcelona, Spain.

²CIBEROBN Fisiopatología de la Obesidad y Nutrición, Instituto de Salud Carlos III, Government of Spain.

³Department of Internal Medicine, Hospital Clínic, IDIBAPS, University of Barcelona, Barcelona, Spain.

⁴Department of Preventive Medicine and Public Health, School of Medicine, University of Navarra, Pamplona, Spain.

⁵IdiSNA, Navarra Institute for Health Research, Pamplona, Spain.

⁶Cardiovascular Risk and Nutrition Research Group (CARIN, Regicor Study Group), IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain.

⁷Department of Epidemiology, Preventive Medicine and Public Health, School of Medicine, University of Valencia, Valencia, Spain.

⁸Human Nutrition Unit, University Hospital of Sant Joan de Reus, Department of Biochemistry and Biotechnology, Faculty of Medicine and Health Sciences, IISPV, Rovira i Virgili University, Reus, Spain.

⁹Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA.

¹⁰Departments of Nutrition and Epidemiology, Harvard School of Public Health, Boston, MA, USA.

*Presenting author

Background and objectives:

Lignans are phenolic compounds found in olive oil, whole grains and seeds, among others. A healthy lifestyle and a healthy diet can reduce the risk of chronic diseases, which are a worldwide burden.

The objective was to prospectively evaluate the association between lignan intake and the incidence of chronic diseases within the PREDIMED study (ISRCTN 35739639), which includes 7,447 participants at high cardiovascular risk that were followed a median of 4,8 years.

Methodology:

Dietary intake of lignans was obtained from yearly food frequency questionnaires using the Phenol-explorer database. Time-dependent Cox proportional hazard models were used to estimate hazard ratios (HR) and 95% confidence intervals (CI) for chronic diseases and mortality according to lignan intake. Statistical analyses were conducted by using SAS software, version 9.3.

Results and conclusions:

After multivariate adjustment and comparing the highest versus the lowest categories, lignan intake was inversely associated with cardiovascular events (HR=0.51, 95% CI 0.30-0.86, P-lineal=0.007), incidence of type 2 diabetes (HR=0.71, 95% CI 0.51-0.98, P-lineal=0.05), and all cause mortality (HR=0.60, 95% CI 0.37-0.97, P-lineal=0.03). In this population, olive oil was the main source of lignans (72%).

Acknowledgements:

CICYT (AGL2016-79113-R) from the Spanish Ministry of Science and Innovation (MICINN), the Generalitat de Catalunya (GC) 2014 SGR 773, the Instituto de Salud Carlos III, ISCIII (CIBEROBN-CB06/03), Institute for Research on Nutrition and Food Safety (INSA-UB), and the European Foundation for Alcohol Research (ERAB) (EA 1324, EA 1514, EA 1515, and EA 1517).